

Immunisation Schedule Queensland

Children and adolescents (10 to 19 years)

July 2018

Always check the Australian Immunisation Register prior to vaccinating

Any additional information about the vaccines included in this schedule should be obtained from the online version of the Australian Immunisation Handbook (AIH).

Age	Disease	Vaccine brand	Reconstitute	Site to administer	Administration	Important notes
Year 7 students (or age equivalent, 11 to 13 years)	Human papillomavirus (HPV)	Gardasil 9	No	Deltoid	IM	<ul style="list-style-type: none"> • 2 doses given at 0 and 6 to 12 months • Immunocompromised adolescents require 3 doses given at 0, 2 and 6 months. • 3 doses for adolescents aged 15 to 19 years given at 0, 2 and 6 months Refer to the ATAGI advice on Gardasil 9
	Diphtheria-tetanus-pertussis	Boostrix	No	Deltoid	IM	<ul style="list-style-type: none"> • 1 dose
Year 10 students (or age equivalent)	Meningococcal ACWY	Menactra	Yes	Deltoid	IM	<ul style="list-style-type: none"> • 1 dose

Additional catch-up vaccines for all children and adolescents aged 10 to 19 years

All 10 to 19 years (with no documented history)	Diphtheria-tetanus-pertussis	Adacel OR Boostrix	No	Deltoid	IM	<ul style="list-style-type: none"> • 3 doses at 0, 1 and 2 months • Minimum interval between each dose is 4 weeks
	Poliomyelitis	IPOL	No	Deltoid	IM	<ul style="list-style-type: none"> • 3 doses at 0, 1 and 2 months • Minimum interval between each dose is 4 weeks
	Measles-mumps-rubella	Priorix OR MMRII	YES	Deltoid	SC or IM SC	<ul style="list-style-type: none"> • 2 doses required at 0 and 1 month • Minimum interval between dose 1 and dose 2 is 4 weeks



Age	Disease	Vaccine brand	Reconstitute	Site to administer	Administration	Important notes
All 10 to 19 years (with no documented history)	Hepatitis B	H-B-VaxII OR Engerix B	No	Deltoid	IM	<p>10 to <11 years and 16 to 19 years (use paediatric vaccine)</p> <ul style="list-style-type: none"> • 3 doses at 0, 1 and 6 months after dose 1 • Minimum interval between dose 1 and dose 3 is 4 months • 11 to 15 years (use adult vaccine) • 2 doses at 0 and 6 months after dose 1 • Minimum interval between dose 1 and dose 2 is 4 months
	Chickenpox	Varilrix OR Varivax	YES	Deltoid	SC	<ul style="list-style-type: none"> • Only 1 dose required for adolescents <14 years • 2 doses required for individuals ≥14 years with a minimum interval of 4 weeks between doses
	Meningococcal ACWY	Menactra	YES	Deltoid	IM	<ul style="list-style-type: none"> • 1 dose for adolescents aged 15 to 19 years <p>Note: If required NeisVacC can be provided for catch up on request</p>

Additional vaccines for Aboriginal and Torres Strait Islander and medically at risk adolescents

Aboriginal and Torres Strait Islanders >15 years	Influenza	As supplied	No	Deltoid	IM	<ul style="list-style-type: none"> • 1 dose every year
People with medical risk factors for invasive pneumococcal disease (IPD)	Pneumococcal	Pneumovax 23	No	Deltoid	IM	Refer to the online version of the AIH for a list of medical risk factors and to determine who is eligible to be vaccinated.

Refer to the online version of The Australian Immunisation Handbook – Table 2.2.2: Recommended needle size, length and angle for administering vaccines