



AGE	ANTIGEN	VACCINE BRAND	COMMENT
CHILDHOOD AND ADOLESCENT			
Birth to <7 days	Hepatitis B	H-B-Vax II Paed or Engerix-B Paed	
6 to 8 weeks	DTPa-hepB-IPV-Hib 13vPCV Rotavirus	Infanrix <i>hexa</i> Prevenar 13 Rotarix	Oral Dose 1: 6 - 14 weeks of age
Aboriginal only	4vMenCV (A,C,W,Y)	Nimenrix	Multiple doses needed # Refer MenACWY Statewide Program
4 months	DTPa-hepB-IPV-Hib 13vPCV Rotavirus	Infanrix <i>hexa</i> Prevenar 13 Rotarix	Oral Dose 2: 10 - 24 weeks of age
Aboriginal only	4vMenCV (A,C,W,Y)	Nimenrix	Multiple doses needed # Refer MenACWY Statewide Program
6 months	DTPa-hepB-IPV-Hib 13vPCV	Infanrix <i>hexa</i> Prevenar 13	
Aboriginal and/or medically at risk	13vPCV	Prevenar 13	
12 months	MMR 13vPCV 4vMenCV (A,C,W,Y)	MMR II or Priorix Prevenar 13 Nimenrix	
Aboriginal only	Hepatitis A	Vaqta	
Pre-term or low birth weight	Hepatitis B	H-B-Vax II Paed or Engerix-B Paed	<32 weeks gestation or <2000g birth weight
13 months to < 5 years	4vMenCV (A,C,W,Y)	Nimenrix	Administer only if 12 month dose not received # Refer MenACWY Statewide Program
18 months	DTPa Hib MMRV	Infanrix or Tripacel Act-Hib or Menitorix Priorix Tetra or ProQuad	Do not administer MMRV as first dose of MMR containing vaccine if < 4 years of age
Aboriginal only	Hepatitis A	Vaqta	
4 years	DTPa-IPV	Quadracel or Infanrix IPV	
All medically at risk children	23vPPV	Pneumovax 23	
Year 7 and 8	HPV dTpa	<i>Gardasil</i> [®] 9 Boostrix or Adacel	Multiple doses needed. Refer Handbook
Year 10 and ages 15 to < 20 years	4vMenCV (A,C,W,Y)	Nimenrix or Menveo	

Persons aged < 20 years can catch up all vaccines above including IPV, VZV and MenC (MenACWY can be used for Men C catch ups)

MenACWY Vaccination Statewide Program Guidance at http://ww2.health.wa.gov.au/Articles/F_1/Immunisation-schedule-and-catch-up-immunisations

ADULT			
≥ 15 years Aboriginal and medically at risk	23vPPV	Pneumovax 23	
≥ 50 years Aboriginal only	23vPPV	Pneumovax 23	
≥ 65 years	23vPPV	Pneumovax 23	
70 to 79 years	Herpes-zoster	Zostavax	
Pregnant Women	dTpa	Boostrix or Adacel	Each pregnancy (recommended 20 - 32 wks)
Birth year ≥ 1966	MMR	Priorix	Give 2 doses if < 2 doses received previously

Adult refugees and humanitarian entrants are eligible to catch up dTpa, Hepatitis B, IPV, MMR and VZV

INFLUENZA			
6 months to <3 years	Influenza	FluQuadri Junior, Fluarix Tetra	Children aged 6 months to <9 years of age require 2 doses (1 month apart) in the first year they receive the vaccine.
3 years to <5 years	Influenza	FluQuadri, Fluarix Tetra	
≥ 5 years Aboriginal and/or medically at risk	Influenza	Afluria Quad, Fluarix Tetra, FluQuadri	
≥ 65 years	Influenza	Fluad	
Pregnant Women	Influenza	Afluria Quad, Fluarix Tetra, FluQuadri	Each pregnancy, any trimester
WA Health Staff	Influenza	Fluarix Tetra, FluQuadri	HCW ≥ 65 years Fluad

Further influenza vaccine information, refer to the advice within the link below:

[2019 NCIRS Fact Sheet](#)

The term Aboriginal is inclusive of Aboriginal and Torres Strait Islander people

Vaccine brands subject to availability. For further information refer to the Australian Immunisation Handbook at the link below:

<https://immunisationhandbook.health.gov.au/>